


Preop Diet

It is recommended by your surgeon and dietitian to follow a low sugar, liquid diet 10 days before surgery. The purpose of this liquid diet is to deplete your glycogen stores, which in turn will make laparoscopic surgery placement easier. Here are examples of liquids you can have during this time period. The nutritional guidelines are as followed:

Consume these liquids in moderation, they contain larger amounts of sugar.

1. **Juice, Gatorade®, or Powerade®** – limit to 2 cups or less each day. (*Try to choose juices lower in sugar such as apple, orange, or grapefruit*)
2. **Regular jello, popsicles, or no sugar added fudgsicles-** 3 or less servings each day
3. **1 cup of milk** (*skim, 1%, Lactaid®, or 2%*) **or 8 oz. Light Yogurt** (*Dannon Light, Yoplait Light or Colombo Light*) **or ½ cup cottage cheese** - 3 or less each day.
4. **Atkins Shakes or Carb Solutions Shakes.** 1-2 a day (These can be found at places like Rite Aid, King Soopers, Walgreens, and Walmart). **DO NOT BUY:** Slim Fast, Ensure, Boost, etc.)



These liquids can be taken in any amount since they do not contain sugar.

1. Sugar-free beverages such as Crystal Light, Wyler's Light, Sugar-free Kool Aid or Sugar-free Tang, diet Snapple, diet sodas, Fruit2O®, Propel®, coffee or tea with sugar substitute, or water.
2. Broth soups(all varieties)
3. Sugar-free popsicles and sugar-free jello

